

Table 8. Rates of perinatal complications as to laboratory diagnoses of cases with HELLP syndrome.

	Perinatal complication exists n (%)	Perinatal complication does not exist n (%)	p value	OR (%95 CI)
Thrombocyte < 70000 (/mm ³)	12(60)	2(40)	0.420	1.4(0.1-14.1)
Thrombocyte > 70000 (/mm ³)	8(40)	3(60)		
AST >400 (IU/l)	7(35)	1(20)	0.520	1.3(0.9-18.8)
AST<400 (IU/ l)	13(65)	4(80)		
ALT >400 (IU/l)	6(30)	1(20)	0.0.659	1.1(0.8-14.7)
ALT <400 (IU/l)	14(70)	4(80)		
LDH >1400 (IU/l)	10(50)	2(40)	0.870	1.5(0.2-11.3)
LDH <1400 (IU/l)	10(50)	3(60)		
Prothrombin Time >14 (sec)	5(25)	1(20)	0.815	0.6(0.3-11.5)
Prothrombin Time <14 (sec)	15(75)	4(80)		
Fibrinogen <200 mg/dl	10(50)	0	0.041	
Fibrinogen >200 mg/dl	10(50)	5(100)		
Creatinin >2 mg/dl	4(20)	0	0.008	
Creatinin <2 mg/dl	16(80)	5(100)		

Values are given as average (x) ± Standard variation (Sx), n % and Odds Ratio (Confidence Interval at 95% in parenthesis)